

UCRL- 93211
PREPRINT

**CIRCULATION COPY
SUBJECT TO RECALL
IN TWO WEEKS**

**RELATIVISTIC CALCULATION OF ATOMIC
N-SHELL IONIZATION BY PROTONS**

Mau Hsiung Chen
Bernd Crasemann

This Paper Was Prepared For Submittal To
Physical Review A

August 14, 1985

Lawrence
Livermore
National
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government thereof, and shall not be used for advertising or product endorsement purposes.

RELATIVISTIC CALCULATION OF ATOMIC N-SHELL IONIZATION
BY PROTONS

Mau Hsiung Chen

Lawrence Livermore National Laboratory

University of California

Livermore, California 94550

and

Bernd Crasemann

Department of Physics and Chemical Physics Institute

University of Oregon

Eugene, Oregon 97403

Relativistic plane-wave Born-approximation calculations of cross sections for N-shell ionization of ^{83}Bi and ^{92}U by protons with incident energies from 0.1 to 1 MeV are reported. The calculations were carried out by using Dirac-Hartree-Slater wave functions. Binding-energy change and Coulomb deflection were taken into account. The relativistic cross sections are compared with values from nonrelativistic Hartree-Slater wave functions to study the effects of relativity.

I. INTRODUCTION

Existing calculations of cross sections for Coulomb ionization of atomic inner shells by proton impact have been carried out primarily in the plane-wave Born approximation (PWBA) with screened hydrogenic wave functions.¹⁻⁴ To go beyond the first Born approximation, the perturbed-stationary-state approach including energy-loss, Coulomb-deflection, binding, polarization and relativistic corrections (ECPSSR) was developed by Brandt and Lapicki.^{5,6} To look into the effect of more realistic wave functions and of ab initio incorporation of relativity, we have performed a series of relativistic plane-wave Born-approximation (RPWBA) calculations of K-, L-, and M-shell ionization cross sections, using Dirac-Hartree-Slater (DHS) wave functions.⁷⁻⁹ These DHS calculations have now been extended to N-subshell ionization cross sections. In this paper, we report on results for N₁₋₇-subshell ionization of ⁸³Bi and ⁹²U by protons and compare these new theoretical cross sections with nonrelativistic Hartree-Slater results in order to study the effects of relativity.

II. THEORETICAL METHOD

In the plane-wave Born approximation (PWBA),¹ the differential cross section for ejection of an electron from a closed atomic n shell by heavy-charged-particle impact is

$$\frac{d\sigma}{dE_f} = \frac{4\pi}{n^2} Z_1^2 e^4 \frac{M_1}{E_1} (2j_n + 1) \int_{q_{min}}^{q_{max}} \frac{dq}{q^3} |F_{fi}(q)|^2. \quad (1)$$

Here, E_f is the kinetic energy of the ejected electron, $\hbar q$ is the momentum transferred to that electron, and Z_1 , M_1 , and E_1 are the charge, mass, and initial kinetic energy of the projectile, respectively; $F_{f1}(q)$ is the relativistic form factor. The exact limits of the momentum transfer^{5,7} were used in the present calculations.

To take into account the effect caused by the presence of the slow charged projectile in the vicinity of the nucleus during the collision, the N-shell binding energy of the united atom (i.e. of the atom with atomic number Z_2+1 in the case of proton impact) was used in these calculations. This increase in binding energy tends to reduce the ionization cross section.

The effect of the Coulomb repulsion between the projectile and the target nucleus on the N-subshell ionization was taken into account by applying a correction factor to the cross section calculated for a straight-line projectile path.^{10,11} The differential cross sections including the Coulomb-deflection correction can be written as^{10,11}

$$\left(\frac{d\sigma_n}{dE_f}\right)^C = \left(\frac{d\sigma_n}{dE_f}\right)^{PWBA} \exp(-\pi dq_0), \quad (2)$$

where

$$q_0 = (U_n + E_f) / v_1. \quad (3)$$

Here, U_n is the binding energy of an n-shell electron, d is the half-distance between the collision partners at closest approach, and v_1 is the initial projectile speed.

A general computer program, written for calculating relativistic PWBA (RPWBA) ionization cross sections with DHS wave

functions, was employed to evaluate the N_1 -subshell Coulomb ionization cross sections. The atomic form factors were calculated with neutral-atom DHS wave functions.^{12,13} The form-factor integrals were computed by two successive fast Fourier transforms.¹⁴ The detailed numerical procedure is described in Ref. 7.

Nonrelativistic Hartree-Slater (HS) wave functions were generated with the same DHS program by multiplying the speed of light by a factor of 1,000 so as to simulate the nonrelativistic limit.^{15,16} The same general program for evaluating relativistic ionization cross sections was then employed to calculate the Hartree-Slater (HS) PWBA cross sections for comparison with the relativistic results. Identical N_1 -subshell binding energies from relativistic theory¹³ were used in both DHS and HS calculations.

III. RESULTS AND DISCUSSION

In Table I we list the N_1 -, N_2 -, and N_3 -subshell ionization cross sections calculated from the DHS model for ^{83}Bi and ^{92}U under proton impact with incident energies between 0.1 and 1 MeV. Results for the $N_{4,5}$ and $N_{6,7}$ subshells are listed in Table II. The theoretical values listed in Tables I and II are relativistic plane-wave Born cross sections (RPWBA) as well as cross sections corrected for binding and Coulomb-deflection effects (RPWBA-BC). The plane-wave Born cross sections from DHS and HS wave functions are compared in Figs. 1 and 2.

The binding correction reduces the N_1 -subshell ionization cross sections by ~15-25% at $E_1=0.1$ MeV and by ~8% at $E_1=1$ MeV. Coulomb deflection leads to a reduction of N_1 -subshell ionization cross sections by as much as a factor of 2 at $E_1=0.1$ MeV but by only 2.8% at $E_1=1$ MeV.

Relativistic effects on the K-, L-, and M-subshell ionization cross sections have previously been found to be quite important.^{7,8,16-18} For the N shell, we find substantial relativistic effects in the lowest two subshells. The N_1 ionization cross section is enhanced by as much as 50% for $_{83}\text{Bi}$ and 80% for $_{92}\text{U}$ if relativity is included. The N_2 ionization cross section is enhanced by 35% for Bi and 45% for U. On the other hand, the ionization cross sections for the outer subshells N_3 through N_7 are affected by only a few percent when relativity is taken into account. The effect of relativity on the total N-shell ionization cross section can therefore be expected to be minimal, since the $N_{6,7}$ -subshell cross sections are one to two orders of magnitude larger than the $N_{1,2}$ cross sections.

ACKNOWLEDGMENTS

We thank William F. Ballhaus, Jr., Director of the NASA Ames Research Center, for permission to use the computational facilities of the Center, and thank the ARC Computational Chemistry and Aerothermodynamics Branch, particularly David M. Cooper, for their hospitality. We gratefully acknowledge the computational assistance of Mei Chi Chen. This work was

performed in part under the auspices of the Department of Energy by the Lawrence Livermore National Laboratory under Contract No. W-7405-ENG-48. In the University of Oregon, this work was supported in part of the Advanced Research Projects Agency of the Department of Defense, monitored by the Air Force Office of Scientific Research under Contract No. F49620-85-C-0040.

References

1. D.H. Madison and E. Merzbacher, in Atomic Inner-Shell Processes, edited by B. Crasemann (Academic, New York, 1985), Vol. 1, p.1.
2. O. Benka and A. Kropf, At. Data Nucl. Data Tables 22, 219 (1978).
3. D.E. Johnson, G. Basbas, and F.D. McDaniel, At. Data Nucl. Data Tables 24, 1 (1979).
4. G.S. Khandelwal, B.H. Choi, and E. Merzbacher, Atomic Data 1, 103 (1969).
5. W. Brandt and G. Lapicki, Phys. Rev. A 23, 1717 (1981).
6. W. Brandt and G. Lapicki, Phys. Rev. A 20, 465 (1979).
7. M.H. Chen, B. Crasemann, and H. Mark, Phys. Rev. A 26, 1243 (1982).
8. M.H. Chen, B. Crasemann, and H. Mark, Phys. Rev. A 27, 2358 (1983).
9. M.H. Chen and B. Crasemann, At. Data Nucl. Data Tables (in press).

10. J. Bang and J.M. Hansteen, K. Dan. Vidensk. Selsk. Mat.-Fys. Medd. 31, No. 13 (1959).
11. G. Basbas, W. Brandt, and R. Laubert, Phys. Rev. A 7 983 (1973).
12. D.A. Liberman, D.T. Cromer, and J.T. Waber, Comp. Phys. Commun. 2 107 (1971).
13. K.-N. Huang, M. Aoyagi, M.H. Chen, B. Crasemann, and H. Mark, At. Data Nucl. Data Tables 18, 243 (1976).
14. J.D. Talman, J. Comput. Phys. 29, 35 (1978).
15. I.P. Grant, B.J. McKenzie, P.H. Norrington, D.F. Mayers, and N.C. Pyper, Comput. Phys. Commun. 21 207 (1980).
16. M.H. Chen, Phys. Rev. A 30, 2082 (1984).
17. T. Mukoyama and L. Sarkadi, Phys. Rev. A 23, 375 (1981).
18. T. Mukoyama and L. Sarkadi, Phys. Rev. A 25, 1411 (1982).

TABLE I. Relativistic plane-wave Born-approximation cross sections (RPWBA) (in barns), calculated from Dirac-Hartree-Slater wave functions for $N_{1,2,3}$ -subshell ionization of Bi and U by protons of energy E_1 (in MeV). Numbers in parentheses indicate powers of 10, e.g., $1.05(3)=1.05 \times 10^3$.

| E_1 | N_1 | | N_2 | | N_3 | |
|-----------|---------|----------|---------|----------|---------|----------|
| | RPWBA | RPWBA-BC | RPWBA | RPWBA-BC | RPWBA | RPWBA-BC |
| 83^{Bi} | | | | | | |
| 0.1 | 1.05(3) | 3.92(2) | 2.71(3) | 9.83(2) | 1.08(4) | 4.51(3) |
| 0.2 | 4.71(3) | 2.47(3) | 1.28(4) | 7.62(3) | 4.16(4) | 2.70(4) |
| 0.3 | 1.42(4) | 9.16(3) | 2.59(4) | 1.81(4) | 7.57(4) | 5.59(4) |
| 0.4 | 2.48(4) | 1.80(4) | 3.83(4) | 2.91(4) | 1.08(5) | 8.48(4) |
| 0.5 | 3.45(4) | 2.68(4) | 4.91(4) | 3.92(4) | 1.36(5) | 1.11(5) |
| 0.6 | 4.27(4) | 3.47(4) | 5.80(4) | 4.79(4) | 1.59(5) | 1.34(5) |
| 0.7 | 4.93(4) | 4.13(4) | 6.50(4) | 5.51(4) | 1.77(5) | 1.53(5) |
| 0.8 | 5.45(4) | 4.67(4) | 7.05(4) | 6.09(4) | 1.91(5) | 1.68(5) |
| 0.9 | 5.85(4) | 5.11(4) | 7.46(4) | 6.55(4) | 2.02(5) | 1.80(5) |
| 1.0 | 6.15(4) | 5.46(4) | 7.78(4) | 6.92(4) | 2.10(5) | 1.89(5) |
| 92^U | | | | | | |
| 0.1 | 3.06(2) | 2.53(2) | 4.38(2) | 1.09(2) | 1.77(3) | 5.47(2) |
| 0.2 | 6.49(2) | 3.34(2) | 1.82(3) | 9.38(2) | 7.60(3) | 4.41(3) |
| 0.3 | 1.78(3) | 1.03(3) | 4.54(3) | 2.89(3) | 1.72(4) | 1.19(4) |
| 0.4 | 4.14(3) | 2.71(3) | 8.09(3) | 5.71(3) | 2.84(4) | 2.14(4) |
| 0.5 | 7.20(3) | 5.14(3) | 1.20(4) | 8.99(3) | 3.99(4) | 3.17(4) |
| 0.6 | 1.05(4) | 7.91(3) | 1.57(4) | 1.23(4) | 5.11(4) | 4.20(4) |
| 0.7 | 1.37(4) | 1.08(4) | 1.93(4) | 1.56(4) | 6.14(4) | 5.17(4) |
| 0.8 | 1.67(4) | 1.36(4) | 2.24(4) | 1.87(4) | 7.06(4) | 6.06(4) |
| 0.9 | 1.94(4) | 1.62(4) | 2.53(4) | 2.14(4) | 7.86(4) | 6.86(4) |
| 1.0 | 2.18(4) | 1.85(4) | 2.76(4) | 2.38(4) | 8.54(4) | 7.55(4) |

TABLE II. Relativistic plane-wave Born-approximation cross sections (RPWBA) (in barns), calculated from Dirac-Hartree-Slater wave functions for $N_{4,5^-}$ and $N_{6,7}$ -subshell ionization of Bi and U by protons of energy E_1 (in MeV).

| E_1 | N_4 | | N_5 | | N_6 | | N_7 | |
|---------------|---------|----------|---------|----------|---------|----------|---------|----------|
| | RPWBA | RPWBA-BC | RPWBA | RPWBA-BC | RPWBA | RPWBA-BC | RPWBA | RPWBA-BC |
| 83Bi | | | | | | | | |
| 0.1 | 6.04(4) | 3.17(4) | 9.76(4) | 5.20(4) | 1.29(6) | 8.07(5) | 1.84(6) | 1.16(6) |
| 0.2 | 1.32(5) | 9.24(4) | 2.29(5) | 1.60(5) | 1.79(6) | 1.36(6) | 2.54(6) | 1.93(6) |
| 0.3 | 1.99(5) | 1.53(5) | 3.49(5) | 2.70(5) | 2.01(6) | 1.63(6) | 2.83(6) | 2.31(6) |
| 0.4 | 2.52(5) | 2.04(5) | 4.37(5) | 3.57(5) | 2.12(6) | 1.78(6) | 2.98(6) | 2.51(6) |
| 0.5 | 2.92(5) | 2.45(5) | 5.01(5) | 4.24(5) | 2.18(6) | 1.87(6) | 3.07(6) | 2.64(6) |
| 0.6 | 3.21(5) | 2.76(5) | 5.47(5) | 4.73(5) | 2.22(6) | 1.93(6) | 3.12(6) | 2.72(6) |
| 0.7 | 3.43(5) | 3.00(5) | 5.79(5) | 5.10(5) | 2.23(6) | 1.97(6) | 3.13(6) | 2.77(6) |
| 0.8 | 3.58(5) | 3.17(5) | 6.02(5) | 5.37(5) | 2.23(6) | 1.99(6) | 3.13(6) | 2.79(6) |
| 0.9 | 3.69(5) | 3.31(5) | 6.19(5) | 5.58(5) | 2.22(6) | 2.00(6) | 3.11(6) | 2.80(6) |
| 1.0 | 3.76(5) | 3.41(5) | 6.28(5) | 5.71(5) | 2.21(6) | 2.00(6) | 3.09(6) | 2.80(6) |
| 92U | | | | | | | | |
| 0.1 | 1.03(4) | 4.06(3) | 1.80(4) | 7.46(3) | 1.64(5) | 8.84(4) | 2.41(5) | 1.32(5) |
| 0.2 | 2.96(4) | 1.92(4) | 4.98(4) | 3.27(4) | 3.02(5) | 2.20(5) | 4.37(5) | 3.22(5) |
| 0.3 | 4.94(4) | 3.66(4) | 8.67(4) | 6.46(4) | 3.99(5) | 3.19(5) | 5.73(5) | 4.61(5) |
| 0.4 | 6.85(4) | 5.42(4) | 1.22(5) | 9.72(4) | 4.72(5) | 3.94(5) | 6.75(5) | 5.66(5) |
| 0.5 | 8.58(4) | 7.05(4) | 1.54(5) | 1.27(5) | 5.29(5) | 4.53(5) | 7.54(5) | 6.48(5) |
| 0.6 | 1.01(5) | 8.53(4) | 1.80(5) | 1.53(5) | 5.73(5) | 4.99(5) | 8.17(5) | 7.14(5) |
| 0.7 | 1.14(5) | 9.83(4) | 2.02(5) | 1.76(5) | 6.10(5) | 5.38(5) | 8.69(5) | 7.69(5) |
| 0.8 | 1.25(5) | 1.10(5) | 2.21(5) | 1.95(5) | 6.40(5) | 5.70(5) | 9.08(5) | 8.12(5) |
| 0.9 | 1.35(5) | 1.20(5) | 2.37(5) | 2.11(5) | 6.63(5) | 5.96(5) | 9.42(5) | 8.49(5) |
| 1.0 | 1.43(5) | 1.28(5) | 2.50(5) | 2.25(5) | 6.82(5) | 6.17(5) | 9.68(5) | 8.78(5) |

Figure Captions

FIG. 1. Cross sections for N_1 -subshell ionization of Bi by proton bombardment, as functions of incident-proton energy. Results of the present relativistic plane-wave Born-approximation calculations from Dirac-Hartree-Slater wave functions (solid curves) are compared with cross sections computed nonrelativistically from Hartree-Slater wave functions (dashed curves).

FIG. 2. Cross sections for N_1 -subshell ionization of U by protons, as functions of incident proton energy. Curves are identified in the caption of Fig. 1.




FIG. 1




FIG. 2